

A p -adic property of Fourier coefficients of modular forms of half integral weight

P. Guerzhoy¹

Notations and Introduction

Denote by M_{2k} (resp. S_{2k}) the space of modular (resp. cusp) forms of weight $2k$ on $SL_2(\mathbb{Z})$. We will write q for $\exp(2\pi i\tau)$, where τ is the variable on the upper-half complex plane. Denote by $M_{k+1/2}^+$ (resp. $S_{k+1/2}^+$) the "+"-subspaces of the spaces of modular (resp. cusp-) forms of half integral weight $k + 1/2$. These subspaces were introduced by Kohnen [5].

Through out the paper we fix an odd prime p .

Let k be an even positive integer.

Definition 1 *We call a pair $(p, 2k)$ singular if $2k \equiv 4, 6, 8, 10$ or $14 \pmod{p-1}$*

Note that each pair of the type $(p, \text{even integer})$ is singular if $p = 3, 5$ or 7 . For each p there exist infinitely many values of k such that the pair $(p, 2k)$ becomes singular. For each k there exist a finite nonempty set of appropriate values of p .

Denote by $\mathbb{C}_p = \hat{\mathbb{Q}}_p$ Tate's field. We fix once and for all an embedding $i_p : \bar{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$. We will not make difference between elements of $\bar{\mathbb{Q}}$ and their images under i_p . The symbol \lim will always denote the limit in \mathbb{C}_p . We write $\sum^{(p)}$ for an infinite sum considered under p -adic topology. Denote by $L_p(s, \chi)$ the p -adic L -function, where χ is a Dirichlet character ([4], p.29-30). We put $\zeta^*(s) = L_p(s, \omega^{1-s})$, where ω is the Teichmüller character.

Following [2] we denote by $H(r, N)$ the generalized class numbers. They coincide with the usual class numbers of binary positive definite quadratic forms when $r = 1$. They are the Fourier coefficients of the unique Eisenstein series

$$\mathcal{H}_{k+1/2} = \zeta(1-2r) + \sum_{N \geq 1} H(k, N)q^N \in M_{k+1/2}^+.$$

One has

$$H(r, N) = L(1-r, \chi) \sum_{d|v} \mu(d)\chi(d)d^{r-1}\sigma_{2r-1}(v/d), \tag{1}$$

where $(-1)^r N = Dv^2$, D is a discriminant of a quadratic field, χ is the Dirichlet character associated with this quadratic field, $\sigma_{2r-1}(n) = \sum_{d|n} d^{2r-1}$ and μ is the Möbius function.

¹Supported by Minerva fellowship

Let $-\Delta$ be the discriminant of the imaginary quadratic field $\mathbb{Q}(\sqrt{-\Delta})$. Let ψ_l , $l \geq 0$ be the theta series associated with the binary quadratic form Q_l , where

$$Q_l(x, y) = \begin{cases} p^{2l} \frac{\Delta}{4} x^2 + y^2 & \text{if } \Delta \equiv 0 \pmod{4} \\ \frac{p^{2l} \Delta + 1}{4} x^2 + xy + y^2 & \text{if } \Delta \equiv -1 \pmod{4}. \end{cases}$$

Put $\psi_l = \sum_{x, y \in \mathbb{Z}} q^{Q_l(x, y)} = \sum_{n \geq 0} b_l(n) q^n$.

Let $f = \sum_{n > 0} a(n) q^n$ be a cusp Hecke eigenform of weight k on $SL_2(\mathbb{Z})$. Denote by $L_2(s, f)$ its symmetric square:

$$L_2(s, f) = \prod_{r: \text{prime}} (1 - \alpha_r^2 r^{-s})^{-1} (1 - \alpha_r \beta_r r^{-s})^{-1} (1 - \beta_r^2 r^{-s})^{-1},$$

where α_r and β_r are complex numbers such that $\alpha_r + \beta_r = r$ and $\alpha_r \beta_r = r^{k-1}$.

Consider Rankin's convolutions

$$D(s, f, \psi_l) = \sum_{n > 0} a(n) b_l(n) n^{-s}$$

The number $D^*(k-1, f, \psi_l) = \pi^{2k-2} L_2(2k-2, f)^{-1} D(k-1, f, \psi_l)$ is algebraic [7]

In the present paper we prove

Theorem 1 *Let $(p, 2k-2)$ be a singular pair. Let χ denote the quadratic Dirichlet character associated with $\mathbb{Q}(\sqrt{-\Delta})$. Then*

$$\begin{aligned} D^*(k-1, f, \psi_0) + (1 - \chi(p) p^{1-k}) \sum_{l > 0}^{(p)} p^{l(2k-3)} D^*(k-1, f, \psi_l) \\ = (1 - \chi(p) p^{1-k}) \frac{(2k-2)!}{2^{2k-3} (k-1) (1 - p^{2k-3}) \zeta(3-2k)} \end{aligned} \tag{2}$$

Remarks

1. It is amusing to notice that the value in the right-hand side of (2) does not depend on the particular choice of the cusp Hecke eigenform f of weight k . The dependence on the choice of the discriminant $-\Delta$ is very slight and explicit. Actually, only the value $\chi(p)$ is involved.

2. The denominators of the numbers $D^*(k-1, f, \psi_l)$ were studied in [8], Theorem 4. Rankin's method was used for this purpose. Even the p -adic convergence of the series (2) does not follow from this result. This convergence is the peculiarity of our singular situation.

We are going to derive theorem 1 from the following

Theorem 2 *Let $(p, 2k)$ be a singular pair. Consider a modular form $\varphi \in M_{k+1/2}^+$. Suppose that $\varphi = \sum_{n \geq 0} c(n) q^n$, $c(n) \in \bar{\mathbb{Q}}$ for all n . Choose N such that $c(N) \neq 0$.*

Then

$$\lim_{r \rightarrow \infty} c(p^r N) = c(0) \frac{L_p(1-k, \chi)}{\zeta^*(1-2k)},$$

where χ is the quadratic character associated with $\mathbb{Q}(\sqrt{(-1)^k N})$.

We prove theorem 2 in Chapter 1. In order to illustrate this theorem, we need to consider modular forms of half integral weight whose Fourier coefficients are "interesting" numbers. Chapter 2 is devoted to theta series. In Chapter 3 we provide a construction which generates another type of half integral weight modular forms. It allows to prove theorem 1. Since this construction seems to us to be of independent interest we will briefly recall it here.

Consider a modular form f of weight k . We suppose that f is a normalized cusp Hecke eigenform. Let F be the Klingen - Eisenstein series associated with f . Since F is a Siegel modular form of degree 2, it has a Fourier-Jacobi expansion ([3], Chapter II): $F = \sum_{m \geq 0} \phi_m(\tau, z) \exp(2\pi i m \tau')$. Here ϕ_m are Jacobi forms of indexes m and the same weight k . One has $\phi_0 = f$. Consider the Jacobi form ϕ_1 . It follows from [3], Theorem 5.4 that ϕ_1 corresponds to a half integral weight modular form φ of weight $k - 1/2$. The form φ belongs to the Kohnen's "+"-space. The Fourier coefficients of the Siegel modular form F were calculated by Böcherer [1] and Mizumoto [7], [8]. These numbers involve special values of Rankin's convolutions of the modular form f with theta series of weight 1. We will apply theorem 2 to the modular form φ . It will yield theorem 1.

Chapter 1

In this chapter we prove theorem 2. First we prepare a few lemmas.

Lemma 1 *Let $(p, 2k)$ be a singular pair. Consider a cusp Hecke eigenform $f = \sum_{n \geq 1} a(n)q^n$ of weight $2k$. Suppose that $a(1) = 1$. Let $K = \mathbb{Q}(a(n)_{n \geq 1})$ be the field extension. Let \mathfrak{P} be a prime ideal in K dividing p . Then \mathfrak{P} divides $a(p)$.*

Remarks

1. K is known to be an algebraic number field.
2. This lemma explains the name "singular". It means that if $(p, 2k)$ is a singular pair, then there are no p -ordinary cusp Hecke eigenforms of weight $2k$, i.e. all the cusp Hecke eigenforms are singular.

Proof of lemma 1.

It is known ([6], Theorem 4.4) that the space S_{2k} possesses a basis over \mathbb{C} which consists of cusp forms with rational integer Fourier coefficients. Let $\varphi_1, \dots, \varphi_t$, where $t = \dim S_{2k}$ and $\varphi_i = \sum_{n \geq 0} b_i(n)q^n$ be such a basis. It follows that there exist algebraic numbers $\alpha_i, \dots, \alpha_t$

such that $f = \sum_i \alpha_i \varphi_i$. It follows from [10], Theorem 7 (see also Remark p.216) that $\lim_{n \rightarrow \infty} b_i(p^n) = 0$ for each i . It yields

$$\lim_{n \rightarrow \infty} a(p^n) = \lim_{n \rightarrow \infty} \sum_{1 \leq i \leq t} \alpha_i b_i(p^n) = 0 \quad (3)$$

If $r \geq 0$ then $a(p^{r+1}) = a(p^r) - p^{2k-1}a(p^{r-1})$, because f is a Hecke eigenform. It follows that $a(p^r) \equiv a(p)^r \pmod{p^{2k-1}}$. Taking in account (3) we obtain the assertion of lemma 1.

Lemma 2 *Let $(p, 2k)$ be a singular pair. Consider a cusp Hecke eigenform $\Phi \in S_{k+1/2}^+$ Let $\Phi = \sum_{n \geq 1} c(n)q^n$ be its Fourier expansion.*

Then $\lim_{r \rightarrow \infty} c(p^r n) = 0$ for each $n > 0$.

Proof.

It is known [5] that one can pick a cusp normalized ($a(1) = 1$) Hecke eigenform f of weight $2k$, $f = \sum_{n \geq 1} a(n)q^n$ such that for $N, n \geq 1$

$$c(n^2 N) = c(N) \sum_{d|n} \mu(d) \left(\frac{N}{d}\right) d^{k-1} a(n/d).$$

In particular we get for $n = p^r$

$$c(p^{2r} N) = c(N) \left(a(p^{2r}) - \left(\frac{N}{p}\right) p^{k-1} a(p^{2r-1}) \right)$$

Combining this formulae with (3) we obtain the assertion of lemma 2.

Our next assertion immediately follows from (1). However we formulate it as a separate lemma.

Lemma 3 *Let χ be the quadratic character associated with $\mathbb{Q}(\sqrt{(-1)^k N})$.*

Then

$$\lim_{r \rightarrow \infty} H(k, p^r N) = \frac{L_p(1-k, \chi)}{1-p^{2k-1}}.$$

Proof of theorem 2.

Consider the basis of the space $M_{k+1/2}^+$ which consists of the finite set of cusp Hecke eigenforms φ_i together with $\mathcal{H}_{k+1/2}$. One has

$$\varphi = \frac{c(0)}{\zeta(1-2k)} \mathcal{H}_{k+1/2} + \sum_i \beta_i \varphi_i$$

with some algebraic coefficients β_i . The assertion of the theorem follows now from lemma 2 and lemma 3.

Remark

It is a well-known estimate that the absolute values of Fourier coefficients of a cusp form of even weight increase slower than those of an Eisenstein series of the same weight. One can consider theorem 2 as a p -adic analogue of this fact. Roughly speaking, consider a modular form $F = \sum_{n \geq 0} c(n)q^n$. Suppose that $F = \mathcal{G} + \Phi$, where $\mathcal{G} = \sum_{n \geq 0} d(n)q^n$ and Φ is a cusp form. Then $\lim_{r \rightarrow \infty} (c(p^r N) - d(p^r N)) = 0$. We have proven such type of statements both in the integral and in the half integral weight cases. The singularity condition is crucial for our argument.

Classically, such type of argument was applied to the Fourier coefficients of theta series. The first illustration of theorem 2 deals with theta series associated with unimodular positive definite quadratic forms.

Chapter 2

Let Q be an unimodular positive definite quadratic form on a lattice Λ of rank $4k$ and $B(x, y)$ the associated bilinear form with $Q(x) = 1/2B(x, x)$. Let $y \in \Lambda$ be such that $Q(y) = 1$. Then by [3], Theorem 7.1 the function

$$\Theta_{Q,y}(\tau, z) = \sum_{x \in \Lambda} q^{Q(x)} \zeta^{B(x,y)} = \sum_{4n \geq r^2} c(4n - r^2) q^n \zeta^r$$

is a Jacobi form of weight $2k$ and index 1 on $SL_2(\mathbb{Z})$. It is known that the function

$$f = \Theta_{Q,y}(\tau, 0) = \sum_{x \in \Lambda} q^{Q(x)} = \sum_{n \geq 0} r_Q(n) q^n$$

belongs to M_{2k} .

Note that $r_Q(n)$ is the number of representations of an integer n by the form Q . $r_Q(0) = 1$. The number

$$c(4n - r^2) = \#\{x \in \Lambda | Q(x) = n, B(x, y) = r\}$$

depends only on $4n - r^2$ by [3], Theorem 2.2. The function

$$\varphi = \sum_{N \geq 0} c(N) q^N$$

belongs to $M_{2k-1/2}^+$. $c(0) = 1$.

In some cases of low weight one can get precise formulae for the numbers $c(N)$ and $r_Q(n)$ (cf. [9], §6 for the integral weight case and [3], p.84-85 for the half integral weight case). One also has the following asymptotics ([9], §6, Cor. 2):

$$r_Q(n) = \frac{4k}{B_{2k}} \sigma_{2k-1}(n) + O(n^k).$$

For the corresponding p -adic statement, we have the proposition below.

Proposition 1 *Let N be a positive integer.*

a. *Suppose that $(p, 2k)$ is a singular pair, and $(p, N) = 1$. Then*

$$(1 - p^{2k-1}) \lim_{r \rightarrow \infty} r_Q(p^r N) = \frac{4k}{B_{2k}} \sigma_{2k-1}(N).$$

b. *Suppose that $(p, 4k - 2)$ is a singular pair. Then*

$$(1 - p^{4k-3}) \lim_{r \rightarrow \infty} c(p^r N) = \frac{L_p(2 - 2k, \chi)}{\zeta(3 - 4k)},$$

where χ is the quadratic character associated with $\mathbb{Q}(\sqrt{-N})$.

Proof.

Write $f = E_{2k} + \Phi$. Here $\Phi = \sum_{n>0} a(n)q^n$ is a cusp form, and $E = 1 + \frac{4k}{B_{2k}} \sum_{n \geq 1} \sigma_{2k-1}(n)q^n$ is the Eisenstein series of weight $2k$. Since $(p, 2k)$ is a singular pair, it follows from [10], Theorem 7 (and loc.cit., Remark, p.216), that $\lim_{r \rightarrow \infty} a(p^r N) = 0$. This yields part **a**. Application of our theorem 2 to the modular form φ implies **b**.

One has to compare our argument with the famous Siegel formulae on theta series. Let us reformulate Siegel's result as an identity of Jacobi forms. For this purpose we briefly recall some notations and definitions from [3]. These notations and definitions will be also useful for us in the next chapter.

Let $k > 2$ be an even number.

Let (Λ_i, Q_i) ($1 \leq i \leq h$) denote the inequivalent unimodular positive definite quadratic forms of rank $2k$ and w_i the number of automorphisms of Q_i . Put $\varepsilon_i = w_i^{-1}/(w_1^{-1} + \dots + w_h^{-1})$. Following [3], Chapter I, we denote by $E_{k,1}$ the Jacobi - Eisenstein series of weight k and index 1. The V_l operator sends a Jacobi form of index m to a Jacobi form of index ml . This operator preserves the weight. Its action on the Fourier expansion coefficients is described by formulae ([3], Theorem 4.2):

$$\phi|V_l = \sum_{n,r} \left(\sum_{a|(n,r,l)} a^{k-1} c(nl/a^2, r/a) \right) q^n \zeta^r.$$

Here $\phi = \sum_{n,r} c(n,r)q^n \zeta^r$ is the Fourier expansion of a Jacobi form ϕ of index m and weight k .

Now Siegel's formulae can be written down as ([3], p.87)

$$\sum_{1 \leq i \leq h} \varepsilon_i \sum_{\substack{y \in \Lambda_i \\ Q_i(y)=m}} \Theta_{Q_i, y}(\tau, z) = (E_{k,1}|V_m)(\tau, z). \quad (4)$$

We are interested in the cases when $m = 0, 1$. If $m = 0$, (4) becomes an identity of modular forms of even weight k . If $m = 1$, (4) becomes an identity of Jacobi forms of index 1 and weight k . Due to the isomorphism established in [3], Chapter II, one can rewrite it as an identity of modular forms of half integral weight $k - 1/2$. In both cases the modular form

which appears in the right hand side is an Eisenstein series. The modular forms which appears in the left hand side of (4), are linear combinations of theta series. We apply our proposition 1 to these theta series. The proposition asserts that for certain p their Fourier coefficients which numbers are divisible by an increasing power of p , become p -adically close to the appropriate coefficients of the Eisenstein series. (See also the remark after the proof of theorem 2.) It accords with (4) since $\sum_{1 \leq i \leq h} \varepsilon_i = 1$.

Chapter 3

This chapter is devoted to the proof of theorem 1.

Let

$$F(Z) = \sum_{T \geq 0} A(T) \exp(2\pi i t r(TZ)) \quad (5)$$

be a Siegel modular form of degree 2 and even weight k . Put $T = \begin{pmatrix} n & r/2 \\ r/2 & m \end{pmatrix}$, where $n, r, m \in \mathbb{Z}$ and $n, m, 4nm - r^2 \geq 0$. Rewrite the Fourier expansion (5) as

$$F(\tau, z, \tau') = \sum_{n, m, 4nm - r^2 \geq 0} A(n, r, m) \exp(2\pi i(n\tau + rz + m\tau')).$$

Lemma 4 *The numbers $A(n, r, 1)$ depend only on $4n - r^2$.*

Define the numbers $c(N)$ by

$$c(N) = \begin{cases} A(n, r, 1) & \text{if there exists a pair } n, r \text{ such that } N = 4n - r^2 \\ 0 & \text{otherwise.} \end{cases}$$

Then the function $\varphi(\tau) = \sum_{N \geq 0} c(N) q^N$ belongs to $M_{k-1/2}^+$.

Proof.

Consider the Fourier-Jacobi expansion of the Siegel modular form F :

$$F(\tau, z, \tau') = \sum_{m \geq 0} \phi_m(\tau, z) \exp(2\pi i m \tau').$$

It follows from [3], Theorem 6.1, that $\phi_m(\tau, z)$ is a Jacobi form of weight k and index m . Consider $\phi_1(\tau, z)$:

$$\phi_1(\tau, z) = \sum_{n, 4n - r^2 \geq 0} A(n, r, 1) \exp(2\pi i(n\tau + rz)).$$

$A(n, r, 1)$ depends only on $4n - r^2$ by [3], Theorem 2.2. The last assertion of the lemma follows from [3], Theorem 5.4.

Let us specialize our consideration to the case when the Siegel modular form is a Klingen - Eisenstein series. The following proposition is a specialization of results obtained in [1], [7], [8].

Proposition 2 Consider $f \in S_k$. Let

$$F(Z) = \sum_{T \geq 0} A(T) \exp(2\pi i t r(TZ))$$

be the Klingen - Eisenstein series associated with f .

Put $T = \begin{pmatrix} n & r/2 \\ r/2 & m \end{pmatrix}$, where $n, r, m \in \mathbb{Z}$ and $n, m, 4nm - r^2 \geq 0$, $\text{g.c.d.}(n, m, r) = 1$.

Suppose that $4nm - r^2 = p^{2\nu} \Delta$, where $-\Delta$ is a fundamental discriminant.

For a positive integer v put

$$\begin{aligned} \Theta_T &= \sum_{x,y} q^{nx^2+rxym+my^2} = \sum_{n \geq 0} b_T(n) q^n, \\ \Theta_T^{(v)} &= \sum_{n \geq 0} b_T(nv^2) q^n. \end{aligned}$$

Consider the algebraic numbers

$$D(T, \mu) = \frac{(k-1)(2\pi)^{2k-2}}{2(2k-2)! L_2(2k-2, f)} D(k-1, f, \Theta_T^{(p^\mu)}).$$

Then

$$A(T) = L(2-k, \chi) \left(D(T, n) + \sum_{0 \leq l < n} p^{(\nu-\mu)(2k-3)} (1 - \chi(p)p^{1-k}) D(T, l) \right).$$

Here χ is the quadratic Dirichlet character associated with $\mathbb{Q}(\sqrt{-\Delta})$, and $L(2-k, \chi)$ is the value at negative integer of the Dirichlet L -function.

We will use proposition 2 in the special case when $m = 1$. In this case $T = \begin{pmatrix} n & r/2 \\ r/2 & 1 \end{pmatrix}$ is the matrix of a quadratic form from the principal class. In what follows we will not make difference between a binary quadratic form and its matrix.

Lemma 5 Suppose that $4n - r^2 = \Delta p^{2\nu}$, $0 \leq \mu \leq \nu$.

There exists a binary quadratic form S with discriminant $-\Delta p^{2\nu-2\mu}$ which belongs to the principal class such that

$$D(T, \mu) = D(S, 0). \tag{6}$$

Proof.

Actually we are going to prove that $\Theta_T^{(p^\mu)} = \Theta_S$. It will yield (6). We claim that if S exists then it belongs to the principal class. If $\Theta_T = \sum_{n \geq 0} b_T(n) q^n$ then $\Theta_S = \Theta_T^{(p^\mu)} = \sum_{n \geq 0} b_T(np^{2\mu}) q^n$. Since T represents 1, $b_T(1) \neq 0$. It yields $b_T(p^{2\mu}) \neq 0$. It means that S represents 1 and our claim follows. The rest of the proof (the existence of S) essentially is contained in [1], p.33. We omit it.

Combining lemma 5 with Proposition 2 we get the explicit formulas for the Fourier coefficients $A(n, r, 1)$ of the Klingen - Eisenstein series F . Lemma 4 allows to regard these numbers as the Fourier coefficients of a modular form of half integral weight. Application of the theorem 2 to these Fourier coefficients completes the proof of theorem 1.

Acknowledgment

It is pleasant for the author to express his gratitude to the Max-Planck-Institut für Mathematik at Bonn for its hospitality and support. The author is very grateful to Professor D. Zagier for a number of useful discussions. The author expresses his gratitude to Professor A. Panchishkin (Institut Fourier, Grenoble) who turned the author's attention to the explicit formulas of Böcherer and Mizumoto.

References

- [1] Böcherer, S. Über gewisse Siegelsche Modulformen zweiten Grades, *Math. Ann.* 261 (1982), 23-41.
- [2] Cohen, H., Sums involving the values at negative integers of L -functions of quadratic characters, *Math. Ann.* 217 (1975), 271-285.
- [3] Eichler, M., Zagier, D., *The theory of Jacobi forms*, Progress in Math. 55, Birkhauser, Boston, 1985.
- [4] Iwasawa, K., *Lectures on p -adic L functions*, Ann. Math. Studies 74, Princeton Univ. Press, 1972.
- [5] Kohlen, W., Modular forms of half-integral weight on $\Gamma_0(4)$, *Math. Ann.* 248 (1980), 249-266.
- [6] Lang, S., *Introduction to modular forms*, Grundlehren der Math. Wissenschaften 222, Springer, 1976.
- [7] Mizumoto, S.-i., Fourier coefficients of generalized Eisenstein series of degree two. I, *Invent. Math.* 65 (1981), 115-135.
- [8] Mizumoto, S.-i., On integrality of certain algebraic numbers associated with modular forms, *Math. Ann.* 265 (1983), 119-135.
- [9] Serre, J.-P., *Cours d'arithmétique*, Presses Universitaires de France, Paris 1970.
- [10] Serre, J.-P., Formes modulaires et fonctions zeta p -adiques, in *Modular functions in one variable*, III, Lect. Notes in Math. 350, 191-268, Springer 1972.